



Editorial Comment: Does Surgical Treatment for Benign Prostate Enlargement (BPE)-Related Bladder Outlet Obstruction (BOO) Benefit Patients with Central Nervous System Diseases? A Systematic Review

Ioannis Charalampous¹, Ioannis Tsikopoulos², Calypso Mitkani³, Michael Samarinas², Yuhong Yuan⁴, Ioannis Vouros¹, Petros Tsafrakidis^{5,6}, Anastasiadis Anastasios⁷, Anastasia Gkotsi⁶, Vasileios Sakalis^{1,6}

¹ Department of Urology, Hippokrateion Hospital of Thessaloniki, 54642 Thessaloniki, Greece; ² Department of Urology, General Hospital of Larisa, 41221 Larisa, Greece; ³ Department of Neurology, Agios Pavlos General Hospital of Thessaloniki, 55132 Thessaloniki, Greece; ⁴ Division of Gastroenterology, Cochrane UGPD Group, McMaster University, Medicine Health Sciences Center, Hamilton, ON L8S 4L8, Canada; ⁵ Germal Oncology Center, Limassol 4108, Cyprus; ⁶ Innovative Surgical and Urological Research Hub (ISUReH), 54250 Thessaloniki, Greece; ⁷ 1st Department of Urology, Aristotele University of Thessaloniki, 54635 Thessaloniki, Greece

J Clin Med. 2024 Sep 30;13(19):5846.

DOI: 10.3390/jcm13195846 | ACCESS: 39407906

Iago Zang Pires¹, Marcio A. Averbek^{2,3}

¹ Divisão de Medicina, Pontifícia Universidade Católica do Rio Grande do Sul - PUCRS, Porto Alegre, RS, Brasil; ² Departamento de Urologia, Hospital São Lucas, Pontifícia Universidade Católica do Rio Grande do Sul - PUCRS, Porto Alegre, RS, Brasil; ³ Coordenador de Neurourologia do Núcleo de Disfunções Miccionais do Hospital Moinhos de Vento, Porto Alegre, RS, Brasil

COMMENT

The risk of bladder outlet obstruction (BOO) related to benign prostate enlargement (BPE) increases with age and concerns all men, even those with central nervous system (CNS) diseases, who can also experience lower urinary tract symptoms (LUTS) secondary to the neurogenic lower urinary tract dysfunction (NLUTD) itself (1). Although prostatic surgery is well known to improve outcomes in BPE-related BOO, its efficacy and safety in patients with CNS diseases generates controversy (2, 3). Charalampous et al. conducted a PRISMA systematic review to evaluate postoperative outcomes after surgery for BPE-Related BOO in men with CNS diseases and NLUTD. The review included 13 studies and involved 1,144 patients stratified in four groups: spinal cord injury, Parkinson's disease, post-cerebrovascular accident, and

multiple system atrophy. Transurethral resection of the prostate (TURP) was the most frequently performed de-obstruction procedure, followed by prostatic artery embolism and open prostatectomy. The primary outcomes pointed to a substantial improvement in symptoms, with a success rate of 81.4% in spinal cord injury, 27.1% in Parkinson's disease, and 66.7% in post-cerebrovascular accident populations. Continence status was assessed in six studies, pointing out a high risk of postoperative urinary incontinence, particularly in patients with multiple system atrophy, 60% of whom developed de novo incontinence symptoms. Perioperative complications, such as urinary infections, were more prevalent in spinal cord injury patients, and higher perioperative mortality rates were observed in post-cerebrovascular accident patients. These results emphasize the possible benefits of prostatic surgery in properly selected neuro-urological patients while highlighting the importance of careful pre-operative assessments to discriminate BPE-related BOO from other etiologies. However, this review only included retrospective studies with a high risk of bias, as well as significant heterogeneity in patient characteristics, diagnosis of BOO, surgery technique, and reported outcomes. Furthermore, most of the studies had a limited number of subjects. Therefore, well-designed prospective studies with standardized inclusion criteria, surgical techniques, and longer follow-up are still needed to determine the real advantage of this intervention in distinct neuro-urological groups.

Correspondence address:**Márcio Averbek, MD, PhD**

Hospital Moinhos de Vento

R. Ramiro Barcelos, 910

Moinhos de Vento

Porto Alegre, RS, 90560-032, Brasil

E-mail: marcioaverbeck@gmail.com

ABBREVIATIONS

BPE = Benign Prostate Enlargement

BOO = Bladder Outlet Obstruction

CNS = Central nervous system

CONFLICT OF INTEREST

None declared.

REFERENCES

1. Ogawa T, Sakakibara R, Kuno S, Ishizuka O, Kitta T, Yoshimura N. Prevalence and treatment of LUTS in patients with Parkinson disease or multiple system atrophy. *Nat Rev Urol.* 2017;14:79-89. doi: 10.1038/nrurol.2016.254.
2. Noordhoff TC, Groen J, Scheepe JR, Blok BFM. Surgical Management of Anatomic Bladder Outlet Obstruction in Males with Neurogenic Bladder Dysfunction: A Systematic Review. *Eur Urol Focus.* 2019;5:875-86. doi: 10.1016/j.euf.2018.02.009.
3. Chang TL, Chen SF, Kuo HC. Surgical outcome of male patients with chronic central nervous system disorders and voiding dysfunction due to bladder outlet obstruction. *Int Urol Nephrol.* 2022;54:2511-9. doi: 10.1007/s11255-022-03285-3.

ARTICLE INFO**Márcio Averbek**<https://orcid.org/0000-0002-8127-7153>

Submitted for publication:
January 01, 2025

Accepted:
January 10, 2025

Published as Ahead of Print:
February 15, 2025